

$\eta_c(2S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.


$\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3638 ± 4 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
3645.0 ± 5.5 ^{+4.9} _{-7.8}	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
3630.8 ± 3.4 ± 1.0	112 ± 24	AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ± 6 ± 8	39 ± 11	CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3639 ± 7	98 ± 52	1 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$
3630 ± 8	164	2 ABE	04G BELL	10.6 $e^+ e^- \rightarrow J/\psi(c\bar{c})$
3622 ± 12	42	2 ABE,K	02 BELL	10.6 $e^+ e^- \rightarrow J/\psi + X$
3594 ± 5		3 EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

² From a fit of the J/ψ recoil mass spectrum. Systematic errors not estimated.

³ Assuming mass of $\psi(2S) = 3686$ MeV.



$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
14 ± 7 OUR AVERAGE					
6.3±12.4±4.0		61	ASNER	04	CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
17.0± 8.3±2.5	112 ± 24		AUBERT	04D	BABR $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<23	90	98 ± 52	⁴ AUBERT	06E	BABR $B^\pm \rightarrow K^\pm X_c \bar{c}$
22 ± 14		121 ± 27	AUBERT	05C	BABR $e^+ e^- \rightarrow J/\psi c \bar{c}$
<55	90	39 ± 11	⁵ CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$
<8.0	95		⁶ EDWARDS	82C	CBAL $e^+ e^- \rightarrow \gamma X$
⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.					
⁵ For a mass value of 3654 ± 6 MeV					
⁶ For a mass value of 3594 ± 5 MeV					

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	
Γ_2 $K\bar{K}\pi$	seen
Γ_3 $p\bar{p}$	
Γ_4 $\gamma\gamma$	seen

$\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	Γ_4
$\Gamma(\gamma\gamma)$	
1.3±0.6 keV	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1.3±0.6 ⁷ ASNER 04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	
⁷ They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.	

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}^2$	$\Gamma_3\Gamma_4/\Gamma^2$
$\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}^2$	
VALUE (units 10^{-8})	
< 5.6	
90 8,9,10 AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 8.0 90 8,9,11 AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$	
<12.0 90 9,11 AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$	
⁸ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.	
⁹ For a total width $\Gamma=5$ MeV.	
¹⁰ For the resonance mass region 3589–3599 MeV/ c^2 .	
¹¹ For the resonance mass region 3575–3660 MeV/ c^2 .	

$\eta_c(2S)$ BRANCHING RATIOS **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_1/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	ABREU	980 DLPH	$e^+ e^- \rightarrow e^+ e^-$ +hadrons	
seen	12 EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$	

 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ
seen	39 ± 11	13 CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$	

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.01	90	LEE	85 CBAL	$\psi' \rightarrow \text{photons}$	
12	For a mass value of 3594 ± 5 MeV				
13	For a mass value of 3654 ± 6 MeV				

 $\eta_c(2S)$ REFERENCES

AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

— OTHER RELATED PAPERS —

BADALIAN	03	PR D67 071901	A.M. Badalian, B.L.G. Bakker	
EICHEN	02	PRL 89 162002	E.J. Eichten, K. Lane, C. Quigg	
ACCIARRI	99T	PL B461 155	M. Acciari <i>et al.</i>	(L3 Collab.)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
PORTER	81	SLAC Summer Inst. 355	F.C. Porter <i>et al.</i>	(CIT, HARV, PRIN+)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)